

1 Developing a task switching training game for children
2 with a rare genetic syndrome linked to intellectual
3 disability.

4 Nigel Robb¹, Annalu Waller², Kate A. Woodcock^{3*}

5 1. Center for Global Communication Strategies, University of Tokyo

6 2. Computing, University of Dundee

7 3. Centre for Applied Psychology, School of Psychology, University of Birmingham

8

9 Corresponding Author:

10 Kate Woodcock

11 papers@katewoodcock.com

12

13

14

15

16

17 [*Nigel Robb and Kate Woodcock were at the School of Psychology, Queen's University](#)
18 [Belfast when the work was conducted.](#)

19

20

21 This article was accepted for publication in Simulation and Gaming on 5th February 2019

22 Robb, N., Waller, A. & Woodcock, K. A. (in press). Developing a task switching training
23 game for children with a rare genetic syndrome linked to intellectual disability. Simulation
24 and Gaming, (Journal Volume Number and Issue Number) pp. xx-xx. Copyright © [2019]
25 (Copyright Holder). DOI: [DOI number]

26 ***Abstract***

27 *Background.* The ability to rapidly switch between tasks is important in a variety of contexts.
28 Training in task switching may be particularly valuable for children with intellectual
29 disability (ID), specifically ID linked to genetic syndromes such as Prader-Willi syndrome
30 (PWS). We have developed a cognitive training game for children with PWS and performed
31 a pilot evaluation of the programme to inform future game development. Here, we describe
32 and critically reflect on the development and pilot evaluation process.

33 *Methods.* Several novel aspects of our approach are highlighted in this paper, including the
34 involvement (in various roles) of children with a rare genetic syndrome (PWS) in the
35 development and evaluation of the software (participatory design) and the development of a
36 matched control, or placebo version of the game for use in the pilot evaluation.

37 *Results.* Children with PWS were capable of contributing to the design and development of a
38 cognitive training game in various roles. In the subsequent pilot evaluation, playing the active
39 version of the game was associated with greater improvement in task switching performance
40 than playing the matched control (placebo) version of the game. However, attrition was an
41 issue during both the design phase and the pilot evaluation.

42 *Conclusions.* The lessons learned from our work have relevance in a wide range of contexts,
43 such as the development of future cognitive training games; the evaluation of serious games
44 in general; and the involvement of end-users with cognitive disabilities and/or rare
45 syndromes in the design and development of software.

46

47 ***Keywords***

48 Game design, participatory design, task switching, executive function, cognitive training,
49 intellectual disabilities

50

51 ***Background***

52 The idea that one could improve one's cognitive skills by playing a game has recently
53 received much attention. Many so-called "brain training" games have been developed and
54 marketed, both by commercial companies and researchers (Rabipour & Raz, 2012). Several
55 of the skills targeted by these games fall under the umbrella of cognitive control, that is,
56 executive process which coordinate and modulate other, more basic cognitive processes.
57 Cognitive control ensures that our various cognitive skills work together in an organised way,
58 which is essential for the completion of complex, goal-directed tasks (Miyake et al. 2000;
59 Miyake & Friedman, 2012). Cognitive control is important for mental and physical health,
60 success at school and work, and quality of life (Diamond, 2013).

61 Cognitive control is also related to a variety of clinically important behaviours. For example,
62 externalising behaviours are associated with poor inhibitory control (Young et al., 2009).
63 Importantly, in children with certain genetic syndromes linked to intellectual disability (ID),
64 deficits in the ability to rapidly switch between tasks (which is a typical cognitive control
65 process) have been associated with a strong resistance to change and preference for routine.
66 In children with the neurodevelopmental disorder Prader-Willi syndrome (PWS), for
67 example, task switching deficits have been causally related to highly negatively impactful
68 behaviours such as temper outbursts (Woodcock, Oliver & Humphreys, 2009). There is also

69 growing evidence supporting a link between task switching and resistance to change – which
70 can also precipitate negatively impactful behavioural problems – in individuals with other
71 neurodevelopmental disorders, such as autism spectrum conditions (Eisenberg et al., 2015;
72 Miller et al., 2015).

73 The possibility of improving control processes such as switching through training games is
74 therefore of much interest and has important applications in clinical populations. However,
75 while the potential of such software as interventions for children with ID is recognised, some
76 current cognitive training programmes may be too advanced for children with ID (Bennett,
77 Holmes & Buckley, 2013). Furthermore, a review published in 2015 includes no studies
78 investigating the effects of training task switching in children with ID (Kirk, Gray, Riby &
79 Cornish, 2015). There is therefore a need to develop and evaluate cognitive control training
80 games which are suitable for children with ID. However, there is currently very little
81 published research which focuses on how such games should be made. Bul et al. (2015)
82 describe the development and user testing of a game designed to improve cognitive control
83 processes in children with Attention Deficit Hyperactivity Disorder (ADHD). While this
84 paper makes an important contribution to our understanding of how user testing of cognitive
85 training games in clinical populations may be carried out, only a brief overview of the
86 “collaborative game development” process is provided. Detailed accounts of methods and
87 results for the development of cognitive training games for clinical populations are lacking.

88 In the present study, we report the design, development, and pilot evaluation of a new task
89 switching training game for children with PWS. Our aim in this paper is to describe the novel
90 aspects of techniques used in the development process, in such a way that the insights
91 obtained – and lessons learned – may benefit those developing and evaluating future
92 cognitive training games. In addition, our findings will be of interest to researchers and
93 designers focused on designing software for people with cognitive disabilities, and those
94 concerned with the development and evaluation of software-based interventions in general.
95 As is important in the design of games, our approach draws on techniques and insights from
96 multiple disciplines. Furthermore, modern software development processes are typically
97 highly flexible and adaptive, in part due to the complexity of software development and the
98 need to respond to change during the development process (Matharu, Mishra, Singh &
99 Upadhyay, 2015). These so-called “agile” approaches are also frequently used in game
100 development (Aleem, Capretz & Ahmed, 2016). As such, the techniques described in this
101 paper were used at various points throughout the development process. However, since these
102 approaches could in principle be applied independently, we present our methods, results, and
103 discussion of those results as two separate studies, each focused on a novel aspect of the
104 development process. Study 1 considers the role of participatory design; i.e., the involvement
105 of end-users in various roles in the development process. Study 2 focuses on the use of active
106 and placebo versions of the software in a pilot evaluation of the prototype game. To our
107 knowledge, our study is the first to use this technique in the evaluation of a cognitive training
108 game.

109 ***Study 1: Participatory design***

110 Understanding the specific needs of the end user of a piece of software (including how they
111 will use the software and in what context) may increase the usefulness and usability of the
112 software (ISO, 2010). One way to achieve such understanding is to involve potential users of
113 the software in the development process; indeed, this is a well-established practice (Bano &
114 Zowghi, 2013). However, more recently, researchers have considered how such involvement
115 may benefit people from traditionally excluded groups, including children with disabilities.
116 For example, researchers have suggested that participation in software design may afford

117 wider benefits to the children involved, such as enjoyment, improved social skills, a sense of
118 empowerment, and increased confidence, in addition to the expected improvements in
119 software quality (Benton & Johnson, 2015). It may therefore be important to involve children
120 with disabilities in the design of cognitive training games. However, as Benton and Johnson
121 (2015) point out, there is still much for researchers to discover about the process of involving
122 children with disabilities in the design of software. In addition, there are various roles which
123 an end user can play in a design process (Benton & Johnson, 2015; Druin, 1999). A meta-
124 analysis of user involvement in the design of games aiming to promote healthy lifestyle
125 behaviours found that certain kinds of user involvement led to games that were less effective
126 than those designed without user involvement (DeSmet et al., 2016). These results were
127 moderated by the role played by users in the design process, and the elements of the game
128 they contributed to. For example, involving users in the aesthetic design of game characters
129 was associated with lower game effectiveness. The authors suggest that design techniques
130 may need to be adapted to suit the users' experience level and cognitive abilities. Combining
131 these two insights shows that there is much to discover about how best to involve children
132 with disabilities (as end-users) in the software development process, including which roles it
133 is most feasible and appropriate for children to take, and the potential benefits and challenges
134 of such user involvement. In the present study, we involved children with PWS in the
135 software development. Here we report on that process with a view to addressing such
136 important questions.

137 ***Methods***

138 ***Participants***

139 A total of eight children (7 – 17 years; 6 female) with PWS and their parents took part in the
140 development process (as is consistent with the individualised nature of responding to
141 feedback ascertained via a participatory design process, children took part in different stages
142 as fitted best into their ongoing lives – leading to differing numbers of participants in each
143 stage). Children and their parents were informed about the research and provided consent in
144 line with procedures as approved by the School of Psychology Research Ethics Committee at
145 Queen's University Belfast. Names used to describe participants in this article are
146 pseudonyms. PWS is a genetic disorder with an estimated lower bound population prevalence
147 rate in the UK of 1:52,000 (Whittington et al., 2001). It is caused by absence of paternally
148 derived genetic material in a specific region (q11-q13) on chromosome 15 (Boer et al., 2002).
149 Since the present study did not seek to inform on phenotypic characteristics of PWS,
150 information on the specific genetic abnormality causing PWS (e.g., paternal deletion) was not
151 sought. Of note, the development of task switching and related cognitive functions has not
152 been examined in individuals with PWS. However, the age range of the present sample
153 corresponds to a period when such cognitive skills and the brain networks underpinning them
154 typically develop greatly (McKenna, Rushe & Woodcock, 2017). This was an important
155 factor contributing to the choice of controlled experimental design in the initial evaluation
156 (reported below).

157 ***Materials and procedure***

158 ***Consultation with children prior to development***

159 Children were consulted at the beginning of the project (i.e., before any design decisions had
160 been made) regarding their preferences and abilities regarding existing games in order to
161 inform on the development of an initial game prototype. Seven children from the overall
162 cohort (age 7 – 15; 6 female) took part. Each child was assisted throughout by a parent. We
163 selected 15 games suitable for children (listed in the Supplementary Materials). This

164 consultation process was previously reported in Robb, Waller & Woodcock (2015). The
165 games were selected to exhibit a variety of design possibilities with respect to the following
166 factors:

167 *Gameplay.* The games selected involved engaging in a variety of activities: solving shape-
168 based puzzles, engaging in intensive multitasking, following simple stories (i.e., less
169 interactivity), collecting items (e.g., coins), controlling animated characters, painting and
170 building virtual objects.

171 *Controls.* Several methods of controlling video games were also presented to the children:
172 using simple touches and gestures on a touchscreen device; using virtual buttons on a
173 touchscreen device (i.e., buttons operated by touching the screen); using a standard computer
174 mouse; using a standard computer keyboard; using motion controls (i.e., using the
175 accelerometer built into most handheld devices to control the game by rotating or otherwise
176 moving the device).

177 *Distribution platforms:* The games selected were played on one of two platforms: handheld
178 devices or computers (either desktop or laptop).

179 Children played each game for five minutes initially, then completed an online questionnaire
180 designed to determine their level of comprehension of each game (example item shown in
181 Figure 1). Children were then given free access to the games for a period of 14 days. The
182 children were free to play whichever games they wanted, although parents were asked to
183 encourage children to consider all the games. At the end of each day, children were asked to
184 complete a short online questionnaire, to determine which games they had played, how long
185 they had played for, and which game was their current favourite (images of the games were
186 displayed; children answered the questions by clicking on the games which they had
187 played/enjoyed). After two weeks, parents and children were asked to complete a more
188 comprehensive online questionnaire. This questionnaire contained detailed questions about
189 the children's favourite and least favourite games, as well as their playing habits and their
190 preferences (example item shown in Figure 2).

191 *Play testing of the initial prototype*

192 After the release of the first playable prototype of the software, children were asked to play
193 the game and provide feedback via online questionnaires (preliminary findings were reported
194 in Robb et al., 2015). Questionnaires incorporated various item types, including Likert-style
195 scales, multiple choice questions, and free-response questions. Questions used simplified
196 language and included images where possible (example item "How much did you enjoy
197 playing the game?"; three-point Likert-style scale with images; see Figure 3). In addition,
198 quiz-style questions were used to establish how well children had understood what they were
199 supposed to do in the game (example item shown in Figure 4).

200 [Figure 1 about here]

201 [Figure 2 about here]

202 [Figure 3 about here]

203 [Figure 4 about here]

204 *Collaborative development*

205 Finally, the lead developer of the game maintained continual contact with the participants'
206 parents (primarily via email) during the entire development process. This communication
207 took several forms. Firstly, parents reported bugs and technical issues as soon as they
208 occurred, allowing these problems to be rectified quickly. Secondly, parents provided

209 descriptions of their child's usage of the game, including how and when they were playing it,
210 what challenges they faced, and what sort of behaviours they displayed. Finally, the lead
211 developer encouraged parents to, in turn, encourage their child to play the game frequently.
212 This ongoing collaborative communication reflects the agile approach to software
213 development used in this project; such agility is widely accepted as an optimal technique in
214 the development of software including video games (Matharu et al., 2015; Aleem et al.
215 2016). This allowed us to adapt the game rapidly in response to feedback, and continually
216 refine the software.

217 ***Results and discussion***

218 Regarding children's preferences and habits in existing games, we found that there was
219 variation in the gameplay preferences of children (Figure 5). When asked to select their
220 favourite game, only one of the gameplay features identified above (intensive multitasking)
221 was not represented. Children generally preferred using simple touch gestures and on-screen
222 buttons to control games, although some children preferred using a keyboard (Figure 6). Less
223 variation was apparent when asked about the platform on which children preferred to play
224 games, with all but one child preferring to play games on a handheld tablet (see also Robb et
225 al., 2015). When asked what they enjoyed about their favourite game, the most popular
226 answer was that the game was easy to play. When asked what they disliked about their least
227 favourite game, the most popular answer was that it was too hard.

228 [Figure 5 about here]

229 [Figure 6 about here]

230 Results from the questionnaire administered to children after playing the first version of the
231 prototype game showed that the core gameplay involved in playing the game was both
232 understood and enjoyed by children. All but one child played the first version of the game for
233 30 minutes before answering the questionnaire; the remaining child only played the game for
234 5 minutes (see also Robb et al., 2015).

235 Through ongoing communication with parents, we were able to determine more specific
236 results regarding the game we were developing. Although the questionnaire showed that
237 children overall understood the gameplay, parents reported that some children were initially
238 confused by the gameplay, which led to some frustration. In addition, parents reported that, at
239 various stages when the gameplay changed, children were also momentarily confused. Some
240 children asked parents questions about what they should do, while other children exhibited
241 frustrated behaviour. Parents also informed us that they believed the first version of the game
242 was interesting and engaging for children, at least for short periods of time. One child
243 expressed an interest in learning more about the player-controlled character (e.g., by asking a
244 parent what the character was feeling). Parents also reported behaviours indicative of children
245 being engaged and/or enjoying themselves (e.g., saying "yes!", when they were successful in
246 the game). Overall, results showed that the game was, as one parent put it, "a good start".
247 Parents noted that they expected more would be required to make the core gameplay more
248 engaging in the long-term.

249 Later in the development process, parents' feedback primarily focused on the challenges
250 faced by children in playing the game. The software had been refined to provide additional
251 features designed to both (1) increase engagement and (2) aid children in understanding the
252 gameplay. However, particularly when changes in the gameplay occurred, parents now began
253 to state that it was increasingly difficult to encourage children to play the game. The
254 challenges faced were of two general types. Firstly, all parents at some point reported that it

255 was challenging to set aside time during the day to focus on the game. Secondly, some
256 parents noted that as children played the game more often, they became less interested in it.

257 Through this participatory design process, involving (1) initial consultation with children
258 with PWS, (2) play testing of an initial prototype, and (3) a collaborative and adaptive
259 approach to refine this prototype, a prototype cognitive training game was developed.

260 The game was implemented as a web-based application optimized to be playable on tablets
261 and mobile phones using simple touch controls. It was developed by one full-stack developer
262 (lead author of this paper) over approximately 12 months. It is difficult to estimate the exact
263 development time, however, as the developer was also a researcher with associated research
264 duties (on the same project), and game development was part time, with hours varying
265 throughout the development period. The developer was experienced in both software
266 engineering/game design and illustration/animation. As such, graphical assets were either
267 created by the developer or obtained from a database of video game assets released under
268 public licences¹. Sound assets were obtained from a similar database of publicly-licenced
269 audio files². The game was developed using Phaser version 2.4.4³. Phaser is a free, open-
270 source, HTML5 game development framework suitable for creating games to be played using
271 desktop or mobile web browsers. It is supported by extensive documentation and examples
272 on the framework's website and online forum, and has a large, active, development and
273 support community. Phaser games may be programmed in either JavaScript or TypeScript; in
274 the current project, JavaScript was used. To ensure a modular, modifiable design, an entity-
275 component system was utilised (West, 2007). Regarding the development environment, a
276 single Windows 7 PC was required. Code was written using the free, open source text editor
277 Atom version 1.x (specific version unknown)⁴. For a local development server, we utilised
278 Mongoose (specific version unknown)⁵. Github⁶ provided version control, and Github Pages⁷
279 was used to host the game directly from the project repository. The project incorporated a
280 backend-as-a-service (BaaS) framework with cloud storage to store persistent data (e.g.,
281 players' accounts). During the work reported here, Parse⁸ (originally developed by Facebook)
282 was used. However, subsequently, Parse was shutdown by Facebook in January 2017, and the
283 framework was open sourced. This required us to migrate to Back4App⁹, which provides a
284 similar BaaS based on the now open source Parse Framework.

285 The core gameplay involved controlling a character to collect items. Although there was
286 variation in children's gameplay preferences, both controlling characters and collecting items
287 were popular among the children. These gameplay mechanics were selected as they provided
288 a simple way to implement task switching demands: the items to be collected were small
289 creatures, although only certain creatures could be collected at any given time. Creatures
290 could be identified in terms of their colour (red or blue) or their shape (cuboid or pyramidal).
291 At some times the target creature was identified by its colour; while at others it was identified
292 by its shape. Children were therefore regularly required to switch between representing the
293 creatures in terms of shape or colour; this provided the core task switching demand of the
294 game (Figure 7, although other additional switching demands were also included. A full

¹ <https://opengameart.org/>

² <https://freesound.org/>

³ <https://phaser.io/>

⁴ <https://atom.io/>

⁵ <https://cesanta.com/>

⁶ <https://github.com/>

⁷ <https://pages.github.com/>

⁸ <https://parseplatform.org/>

⁹ <https://www.back4app.com/>

295 description of the gameplay and the rationale for including specific features is provided in
296 Supplementary Materials (Table S1).

297 By consulting with children before any design decisions had been made, we were able to
298 select a core gameplay mechanic that children understood and were able to use. This suggests
299 that, for certain decisions in a software development process where the end users may have
300 special requirements due to disabilities, participatory design may be an important factor in
301 informing key design decisions. Previous research has identified several roles that children
302 can play in the design process, such as tester, informant, and partner (Druin, 1999). Our
303 findings show that children with intellectual disabilities can play both tester and informant
304 roles (an informant role is ongoing and includes being consulted at an early stage to provide
305 input to initial design decisions) in the design of video games. This provides evidence that
306 future participatory design practices can build on. Although there is now a growing body of
307 research focused on participatory design with children with disabilities, most of this research
308 involves children with autism. Children with other developmental disabilities, particularly
309 rare genetic syndromes such as PWS, are much less likely to be involved in participatory
310 design (Benton & Johnson, 2015; Börjesson, Barendregt, Eriksson & Torgersson, 2015). Our
311 work shows that, by using simplified information and visual elements, and with parental
312 support, such children can successfully inform the development of video games. Future work
313 should expand upon this to include children with a wider range of disabilities in the design of
314 software. However, we also note that, here, children did not contribute to all design decisions.
315 This is of course inevitable: when designing a game to train task switching, at least some of
316 the requirements (e.g., that the player is required to perform task switches) are known in
317 advance. We recommend that future participatory designers reflect on the usefulness of
318 children's contributions before engaging in participatory design and focus children's input on
319 requirements that they can reasonably be expected to contribute to. In this regard the
320 informant role (Druin, 1999), in which children are consulted on an ongoing basis, but only
321 where the designers believe they can make a useful contribution, is perhaps most appropriate
322 in the design of educational and healthcare technology, where there are specific, known
323 requirements that the software must have in order to be effective.

324 ***Study 2: Initial evaluation of the game as including an appropriate*** 325 ***gameplay mechanic for training task switching***

326 Although many studies have been conducted to evaluate the effects of cognitive training
327 programmes, recent reviews and meta-analyses have highlighted the urgent need to improve
328 the design and implementation of evaluation studies (Simons et al., 2016). For example, one
329 issue with cognitive training studies is the difficulty of ensuring a suitably matched control
330 group. Previous studies have used a range of control conditions, including passive controls
331 (i.e., the participants engage in no training), and active controls in which participants may
332 engage in a non-computerised cognitive task (e.g., doing crossword puzzles) or some other
333 kind of game (e.g., playing Tetris) (Simons et al., 2016). Simons et al. (2016) suggest that,
334 unlike in drug trials, for example, it is impossible for participants to be blind to their group-
335 allocation in a cognitive training trial; the participants in the placebo group will be aware that
336 they are not using the cognitive training programme. However, it may be possible to achieve
337 this by creating a version of the cognitive training game which has key features (i.e., features
338 expected to target the cognitive processes being changed) either removed or modulated, but is
339 otherwise identical to the training version. Essentially, this would amount to creating a
340 placebo version of the game software, which could therefore facilitate double-blind placebo-
341 controlled trials of cognitive training games.

342 There are substantial costs involved in game development, and there have been mixed results
343 around the efficacy of game-based cognitive training (e.g. see Karbach & Unger, 2014). A
344 full discussion of the potential merits of computer games for cognitive training is beyond the
345 scope of the present paper. However, it is clear that if pursuing a cognitive training goal with
346 computer game design, evidence that the gameplay mechanic is capable of exerting an
347 appropriate (beneficial) influence on the cognitive process that is the target for the training, is
348 important. Furthermore, obtaining this evidence early in the game development process
349 would ensure that resources can be directed along promising lines.

350 In the present study we therefore aimed to develop a matched placebo version of the early
351 stage cognitive training game that was the result of the participatory design process (study 1).
352 The matched placebo controlled for the whole gameplay experience but did not contain the
353 gameplay features that were designed to place demands on task switching (the active
354 features). Furthermore, we aimed to evaluate the capacity of the active features to influence
355 task switching beneficially in a placebo controlled cross-over design. If the active features
356 benefited task switching ability, this would provide evidence that the early stage game
357 developed via the participatory design process, is suitable for further development.

358 ***Methods***

359 ***Development of the placebo version of the game***

360 During development of the game, features were implemented in the game code in a modular
361 fashion, so that the implementation of each specific feature was, as much as possible,
362 separated from that of other features. It was therefore a straightforward process for the
363 research and development team to create active and matched placebo versions of the game.
364 Specifically, the placebo version of the game did not include any switching demands other
365 than the core switching demand (Figure 7), while the active version of the game included
366 unexpected events which required the player to perform additional task switches (described
367 fully in Supplemental Material, Tables S1 and S2). The active game also featured a difficulty
368 adjustment system which provided increasing challenge over time (i.e., both within
369 individual games and across multiple games) as the player performed better in the game. The
370 placebo game provided a simple difficulty adjustment system which provided increasing
371 challenge within individual games but did not adapt to players' performance over multiple
372 games (i.e., each new game began at the easiest setting, regardless of a player's performance
373 in previous games). The details of these difficulty adjustment systems are explained in
374 Supplemental Material (Table S1).

375 ***Evaluation of the appropriateness of the active game features for training task*** 376 ***switching***

377 ***Participants and design***

378 All 8 individuals with PWS who took part in the design process (9-17 years) were invited to
379 participate in the placebo-controlled, cross-over experiment (see Figure 8). Participants were
380 randomly allocated into one of two groups defining whether they played the active version of
381 the game or a corresponding placebo version. Three participants dropped out before
382 commencement of the test due to other demands for the family at the time, which prevented
383 the time required being available. Five participants began taking part and were asked to play
384 the game as much as possible over a four-week period.

385 [Figure 8 about here]

386 **Measures**

387 Four computer-based neurocognitive task switching tests were administered to index
388 cognitive skill in task switching. The tests were based on the work of Miyake et al. (2000);
389 and adaptations that have been employed with children (e.g., Lehto et al., 2006). The tests are
390 described in detail in the Supplementary Materials. In brief, the tests each presented
391 participants with visual stimuli that needed to be identified as belonging to one of two values
392 of two possible dimensions (e.g. a male or female person in a gender dimension; or a young
393 or old person in an age dimension). Trials where the relevant dimension was different to that
394 in the previous trial demanded a task switch (switch trials), whereas trials where the relevant
395 dimension was the same as that in the previous trial did not demand such as task switch.

396 Importantly, assessing task switching (and more broadly executive functioning) is
397 particularly challenging in individuals with intellectual disabilities (Bevins & Hulse, 2016)
398 because tests of such processes necessarily make demands on a range of lower level cognitive
399 processes that may be selectively impaired. The four tests used here were specifically
400 designed to overcome this challenge by taking participants through a graded practice
401 procedure, which forced participants to demonstrate acceptable ability in all non-switching
402 cognitive processes involved in the test, before they were permitted to continue to the part of
403 the test that assessed task switching.

404 Outcome variables were calculated after screening for outlying trials on an individual
405 participant basis. Switch time was the mean reaction time (in milliseconds) in switch trials;
406 and switch error was the proportion of incorrect switch trials. Composite switching outcome
407 variables were calculated across all tasks completed at every relevant time point. This
408 allowed the simple means of switch time and switch error variables, across all relevant tasks,
409 to constitute composite outcomes that were comparable over time, despite reaction times and
410 error rates not being directly comparable across tests.

411 **Procedure**

412 The switching tests were completed at home via the internet, under the supervision of a
413 parent on four occasions, twice before engagement with placebo or active versions of the
414 game commenced, once following phase 1 of gameplay, and once following phase 2 of
415 gameplay (see Figure 8). Importantly, administration of the switching tests twice before
416 phase 1 of gameplay provided an index of expected improvement in switching test
417 performance driven purely by prior practise with the tests. A brief parent report and self-
418 report questionnaire on behavioural indicators of impaired switching; and on experience
419 during gameplay were also administered via online forms. These questionnaires were
420 pertinent to study goals wider than those described here and so are not discussed further.

421 **Results and discussion**

422 The duration between assessment time points varied for some participants (see Table 1)
423 because for some, daily life disrupted the opportunity to dedicate time to the training, so time
424 elapsed between when an assessment was completed and when training was engaged with.
425 Training time was always accrued primarily during the four weeks preceding the assessment
426 that followed the corresponding training phase.

427 Of the five participants who began taking part, two had been randomly allocated to receive
428 the active training first (Pseudonyms Mary and Jess), whilst the others had been randomly
429 allocated to receive the placebo training first. However, a technical error meant that Mary
430 actually received the placebo training first. As illustrated in Table 1, only one of the four
431 participants who received the placebo training first (Sarah) continued to complete the active
432 training phase.

433 Initially, participants and caregivers were blinded to which type of training the child was
434 completing. However, since those who began with the placebo training lacked motivation to
435 continue, parents of children who began with the placebo training were told about their
436 child's training allocation at the end of phase 1. Two participants finished both training
437 phases, one completed the placebo training first, and the other (Jess), completed the active
438 training first.

439 As expected, all participants demonstrated an improvement in switching linked to practise
440 with the switching tests. Relative to these practice effects however, placebo training was
441 associated with consistently less improvement in performance across all participants. On the
442 other hand, active training (specifically, engagement for at most 2 hours 45 minutes) was
443 associated with more improvement in performance relative to practise for both participants
444 who completed such training (Table 1). Thus, active training did appear capable of improving
445 task switching performance outside of the training environment.

446 [Table 1 about here]

447 Participants' lack of motivation to engage with the placebo training, which led most to drop
448 out before the active training phase, has important implications for the concept of using a
449 non-active version of a cognitive training game as a placebo. If engagement with the placebo
450 game cannot be maintained during the training period, a randomised controlled trial would
451 not be capable of differentiating effects of active training from repeated engagement with any
452 computer-based activity. Furthermore, as evidenced by our results, a cross-over design would
453 be problematic.

454 Given the present early stage of game development, in removing active components from the
455 game to create the placebo version, the only way in which the game adapted to players'
456 ongoing performance was also removed (in the active game, switching demands increased
457 with ongoing play). A similar non-adaptive placebo control approach has been applied
458 previously in systematic evaluations of computer based cognitive training programmes
459 (Spencer-Smith & Klingberg, 2015). However, such previous evaluations have in the most
460 part used a face to face trainer-student set up to provide an external motivator for
461 engagement. On the other hand, an appropriate level of challenge makes an important
462 contribution to intrinsic motivation (Abuhamdeh & Csikszentmihalyi, 2012). The present
463 results highlight the importance of such a source of intrinsic motivation and suggest that
464 future work should aim to provide appropriate challenge in placebo versions of cognitive
465 training games. Indeed, work designing a computer game for training in mathematics has
466 begun to distinguish between how the overall game adapts to challenge players, and how the
467 mathematical content adapts (Mees, Habgood, Jay, & Howard-Jones, 2017).

468 Despite the challenges with placebo control, the gains in performance on the neurocognitive
469 switching tests mediated by engagement with the active training provide important evidence
470 to support the basic game dynamic as a core component of a task switching training
471 programme for people with PWS. Other training programmes that have been linked to
472 beneficial cognitive outcomes usually comprise at least 12 training hours (Spencer-Smith &
473 Klingberg, 2015). The present prototype game lacked scope to encourage play for longer
474 periods of time. Importantly however, demonstrating such a beneficial effect on task
475 switching at the present early stage of game development provides a strong basis for further
476 development of a game based around the present gameplay mechanic.

477 **Conclusion**

478 We developed a prototype game for training task switching in children with PWS, which
479 appears to provide an appropriate foundation for further development of a task switching
480 training computer game for this population. Participatory design allowed a prototype to be
481 created that engaged children for short periods of time. Furthermore, it allowed some
482 important limits to usability to be identified, and the software to be refined to overcome
483 these. However, it is also quite likely that, over a longer period of time, user-involvement in
484 the design process led to fatigue. This is shown by (1) how as development progressed, it
485 became more challenging to encourage participants to use the software, and (2) the levels of
486 attrition experienced in the evaluation of the prototype game. Regarding the former point,
487 research on participatory design has recently begun to focus on the potential benefits for the
488 participants themselves rather than benefits in software quality (Benton & Johnson, 2015).
489 Here we note that there may be conflicts between these two motivations. For example,
490 towards the end of the development process, we sought to obtain feedback from children
491 regarding the usability of the latest version of the software. Even though children did enjoy
492 playing the game, spontaneous, free play is not the same as being asked to play for a specific
493 amount of time then provide feedback (e.g., by completing a questionnaire or verbalising
494 their thoughts to parents). Therefore, it seems likely that children's enjoyment may
495 potentially be at odds with researchers' and developers' need to obtain useful feedback. Of
496 course, in these cases, the voluntary wishes and enjoyment of the children should be put first
497 (as it always was in the project reported here). To address this, participatory designers may
498 need to recruit a larger cohort of participants, thus recognising that some children may not
499 want to be involved in every stage. However, this of course presents a unique challenge when
500 the participants are of interest because they have been diagnosed with a rare syndrome such
501 as PWS.

502 Although the attrition observed following engagement with the placebo training was an
503 important finding of the present study, the attrition post recruitment but before any training
504 had begun was a limitation. This attrition reflected the typical busy lives of families and is
505 important to consider with respect to future trial designs requiring substantial time input from
506 participants as they engage with a cognitive training programme. A related limitation was the
507 variation in time to complete the training phases in the evaluation of the prototype. Periods
508 longer than those planned lapsed between some assessment time points for the two
509 participants who completed both training phases, because of the need to adapt the procedure
510 around participants' lives. It is important to bear this limitation in mind going forward when
511 thinking about how best to encourage regular engagement with a cognitive training game. It
512 may be for example, that games with short chunks of gameplay, which could be completed
513 flexibly around other activities, would be well suited to meet this need. Indeed, our ongoing
514 development of the prototype game described here encompasses such a structure.

515 **References**

- 516 Abuhamdeh, S., & Csikszentmihalyi, M. (2012). The importance of challenge for the
517 enjoyment of intrinsically motivated, goal-directed activities. *Personality and Social*
518 *Psychology Bulletin*, 38(3), 317-330. doi:10.1177/0146167211427147
- 519 Aleem, S., Capretz, L. F., & Ahmed, F. (2016). Game development software engineering
520 process life cycle: a systematic review. *Journal of Software Engineering Research and*
521 *Development*, 4(1), 6. doi: 10.1186/s40411-016-0032-7

- 522 Bano, M., & Zowghi, D. (2013, April). User involvement in software development and
523 system success: a systematic literature review. In *Proceedings of the 17th International*
524 *Conference on Evaluation and Assessment in Software Engineering* (pp. 125-130). ACM.
525 doi: 10.1145/2460999.2461017
- 526 Bennett, S. J., Holmes, J., & Buckley, S. (2013). Computerized memory training leads to
527 sustained improvement in visuospatial short-term memory skills in children with Down
528 syndrome. *American journal on intellectual and developmental disabilities*, 118(3), 179-192.
529 doi: 10.1352/1944-7558-118.3.179
- 530 Benton, L., & Johnson, H. (2015). Widening participation in technology design: A review of
531 the involvement of children with special educational needs and disabilities. *International*
532 *Journal of Child-Computer Interaction*, 3, 23-40. doi: 10.1016/j.ijcci.2015.07.001
- 533 Bevens, S., & Hurse, E. (2016). The assessment of executive functioning in people with
534 intellectual disabilities: an exploratory analysis. *British Journal of Learning Disabilities*,
535 44(2), 87-94. doi:10.1111/bld.12112
- 536 Bisoglio, J., Michaels, T. I., Mervis, J. E., & Ashinoff, B. K. (2014). Cognitive enhancement
537 through action video game training: great expectations require greater evidence. *Frontiers in*
538 *psychology*, 5. doi: 10.3389/fpsyg.2014.00136
- 539 Boer H., Holland A. J., Whittington J. E., Butler J., Webb T. & Clarke D. (2002) Psychotic
540 illness in people with Prader-Willi syndrome due to chromosome 15 maternal uniparental
541 disomy. *The Lancet* 359, 135–6.
- 542 Börjesson, P., Barendregt, W., Eriksson, E., & Torgersson, O. (2015, June). Designing
543 technology for and with developmentally diverse children: a systematic literature review. In
544 *Proceedings of the 14th international conference on interaction design and children* (pp. 79-
545 88). ACM. doi: 10.1145/2771839.2771848
- 546 DeSmet, A., Thompson, D., Baranowski, T., Palmeira, A., Verloigne, M., & De
547 Bourdeaudhuij, I. (2016). Is participatory design associated with the effectiveness of serious
548 digital games for healthy lifestyle promotion? A meta-analysis. *Journal of medical Internet*
549 *research*, 18(4). doi: 10.2196/jmir.4444
- 550 Diamond, A. (2013). Executive functions. *Annual review of psychology*, 64, 135-168. doi:
551 10.1146/annurev-psych-113011-143750
- 552 Dimitropoulos, A., Blackford, J., Walden, T., & Thompson, T. (2006). Compulsive behavior
553 in Prader-Willi syndrome: Examining severity in early childhood. *Research in*
554 *Developmental Disabilities*, 27(2), 190-202. doi:10.1016/j.ridd.2005.01.002
- 555 Druin, A. (1999). The role of children in the design of new technology. *Behaviour and*
556 *Information Technology*, 21(1), 1-25.
- 557 Eisenberg, I. W., Wallace, G. L., Kenworthy, L., Gotts, S. J., & Martin, A. (2015). Insistence
558 on sameness relates to increased covariance of gray matter structure in autism spectrum
559 disorder. *Molecular Autism*, 6, 12. doi:10.1186/s13229-015-0047-7
- 560 Engeser, S., & Rheinberg, F. (2008). Flow, performance and moderators of challenge-skill
561 balance. *Motivation and Emotion*, 32(3), 158-172. doi:10.1007/s11031-008-9102-4
- 562 Gee, J. P. (2007). *What video games have to teach us about learning and literacy*. New York,
563 NY: Palgrave MacMillan.

- 564 Gentile, D. A., & Gentile, J. R. (2008). Violent video games as exemplary teachers: A
565 conceptual analysis. *Journal of Youth and Adolescence*, 37(2), 127-141. doi: 10.1007/s10964-
566 007-9206-2
- 567 Gottlieb, J., Lopes, M., & Oudeyer, P. Y. (2017). Motivated cognition: neural and
568 computational mechanisms of curiosity, attention, and intrinsic motivation. In S. I. Kim, J.
569 Reeve, & M. Bong (Eds.), *Recent Developments in Neuroscience Research on Human*
570 *Motivation* (Vol. 19, pp. 149-172). Bingley: Emerald Group Publishing Ltd.
- 571 Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). Behavior rating inventory
572 of executive function. *Child Neuropsychology*, 3, 235–238.
- 573 Gioia, G. A., Isquith, P. K., Kenworthy, L. & Barton, R. M. (2002) Profiles of everyday
574 executive function in acquired and developmental disorders, *Child Neuropsychology*, 8:2,
575 121-137, DOI: 10.1076/chin.8.2.121.8727
- 576 Gioia, G. A., Isquith, P. K., Retzlaff, P. D. & Espy, K. A. (2002) Confirmatory factor
577 analysis of the Behavior Rating Inventory of Executive Function (BRIEF) in a clinical
578 sample, *Child Neuropsychology*, 8:4, 249-257. doi: 10.1076/chin.8.4.249.13513
- 579 Green, C. S., & Bavelier, D. (2008). Exercising your brain: A review of human brain
580 plasticity and training-induced learning. *Psychology and Aging*, 23(4), 692–701. doi:
581 10.1037/a0014345
- 582 Hartley, S. L. & MacLean, W. E. (2006). A review of the reliability and validity of Likert-
583 type scales for people with intellectual disability. *Journal of Intellectual Disability Research*,
584 50(11), 813-827. doi: 10.1111/j.1365-2788.2006.00844.x
- 585 ISO. (2010) *ISO 9241-210: Ergonomics of human-system interaction – Part 210: Human-*
586 *centred design for interactive systems*. Geneva, Switzerland: International Organization for
587 Standardization.
- 588 Jurado, M. B., & Rosselli, M. (2007). The elusive nature of executive functions: a review of
589 our current understanding. *Neuropsychology review*, 17(3), 213-233. doi: 10.1007/s11065-
590 007-9040-z
- 591 Kagohara, D. M., van der Meer, L., Ramdoss, S., O'Reilly, M. F., Lancioni, G. E., Davis, T.
592 N., ... & Green, V. A. (2013). Using iPods® and iPads® in teaching programs for individuals
593 with developmental disabilities: A systematic review. *Research in developmental disabilities*,
594 34(1), 147-156. doi: 10.1016/j.ridd.2012.07.027
- 595 Karbach, J., & Unger, K. (2014). Executive control training from middle childhood to
596 adolescence. *Frontiers in Psychology*, 5, 14. doi:10.3389/fpsyg.2014.00390
- 597 Kirk, H. E., Gray, K., Riby, D. M., & Cornish, K. M. (2015). Cognitive training as a
598 resolution for early executive function difficulties in children with intellectual disabilities.
599 *Research in developmental disabilities*, 38, 145-160. doi: 10.1016/j.ridd.2014.12.026
- 600 Lehto, J. E., Juuja, P., Kooistra, L., Pulkkinen, L., Juujärvi, P., Kooistra, L., & Pulkkinen, L.
601 (2003). Dimensions of executive functioning : Evidence from children. *British Journal of*
602 *Developmental Psychology*, 21(1), 59–80. <https://doi.org/10.1348/026151003321164627>
- 603 Matharu, G. S., Mishra, A., Singh, H., & Upadhyay, P. (2015). Empirical study of agile
604 software development methodologies: A comparative analysis. *ACM SIGSOFT Software*
605 *Engineering Notes*, 40(1), 1-6. doi: 10.1145/2693208.2693233
- 606 McKenna, R., Rushe, T., & Woodcock, K. A. (2017). Informing the structure of executive
607 function in children: A meta-analysis of functional neuroimaging data. *Frontiers in Human*

608 Neuroscience, 11, 17. Retrieved from <Go to ISI>://WOS:000401589100001.
609 doi:10.3389/fnhum.2017.00154

610 Mees, M., Habgood, J., Jay, T. & Howard-Jones, P. (2017). Researching adaptivity for
611 individual differences in numeracy games. *CHI PLAY'17 Extended Abstracts, Oct. 15–18*,
612 Amsterdam, NL. doi: 10.1145/3130859.3131315

613 Mellecker, R., Lyons, E. J., & Baranowski, T. (2013). Disentangling fun and enjoyment in
614 exergames using an expanded design, play, experience framework: A narrative review.
615 *Games for health*, 2(3), 142-149. doi: 10.1089/g4h.2013.0022

616 Miller, H. L., Ragozzino, M. E., Cook, E. H., Sweeney, J. A., & Mosconi, M. W. (2015).
617 Cognitive set shifting deficits and their relationship to repetitive behaviors in autism
618 spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(3), 805-815.
619 doi:10.1007/s10803-014-2244-1

620 Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D.
621 (2000). The unity and diversity of executive functions and their contributions to complex
622 “Frontal Lobe” tasks: a latent variable analysis. *Cognitive Psychology*, 41(1), 49–100.
623 <https://doi.org/10.1006/cogp.1999.0734>

624 Perkins, D. N., & Salomon, G. (2012). Knowledge to go: A motivational and dispositional
625 view of transfer. *Educational Psychologist*, 47(3), 248-258.
626 doi:10.1080/00461520.2012.693354

627 Pugh, K. J., & Bergin, D. A. (2006). Motivational influences on transfer. *Educational*
628 *Psychologist*, 41(3), 147-160. doi:10.1207/s15326985ep4103_2

629 Rheinberg, F. (2008). Intrinsic motivation and flow-experience. *Motivation and action*,
630 323348. doi: 10.1017/CBO9780511499821.014

631 Robb, N., Waller, A. & Woodcock, K. A. (2015). The development of TASTER, a cognitive
632 training game using human-centered design, tailored for children with global and specific
633 cognitive impairments. *2015 IEEE 7th International Conference on Games and Virtual*
634 *Worlds for Serious Applications (Vs-Games)*, 208-209. Retrieved from <Go to
635 ISI>://WOS:000380426500031.

636 Simons, D. J., Boot, W. R., Charness, N., Gathercole, S. E., Chabris, C. F., Hambrick, D. Z.,
637 & Stine-Morrow, E. A. (2016). Do “brain-training” programs work? *Psychological Science in*
638 *the Public Interest*, 17(3), 103-186. doi: 10.1177/1529100616661983

639 Spencer-Smith, M., & Klingberg, T. (2015). Benefits of a working memory training program
640 for inattention in daily life: A systematic review and meta-analysis. *Plos One*, 10(3), 18.
641 doi:10.1371/journal.pone.0119522

642 Tunnicliffe, P. L., Woodcock, K. A., Bull, L.E, Penhallow, J. & Oliver, C. (2014). Temper
643 outbursts in Prader-Willi syndrome: Causes, behavioural and emotional sequence and
644 responses by carers. *Journal of Intellectual Disability Research*. 58, 134-150. doi:
645 10.1111/jir.12010

646 Weibel, D. & Wissmath, B. (2011). Immersion in computer games: The role of spatial
647 presence and flow. *International Journal of Computer Games Technology*, 282345,
648 doi:10.1155/2011/282345

649 Whittington, J. E., Holland, A. J., Webb, T., Butler, J., Clarke, D. & Boer, H. (2001).
650 Population prevalence and estimated birth incidence and mortality rate for people with
651 Prader-Willi syndrome in one UK Health Region. *Journal of Medical Genetics*, 28, 702-708.

- 652 West, M. (2007). Refactoring game entities with components. Retrieved from
653 [http://cowboyprogramming.com/2007/01/05/evolve-your-heirachy./](http://cowboyprogramming.com/2007/01/05/evolve-your-heirachy/)
- 654 Woodcock, K. A., Oliver, C., & Humphreys, G. W. (2009). Task-switching deficits and
655 repetitive behaviour in genetic neurodevelopmental disorders: data from children with
656 Prader–Willi syndrome chromosome 15 q11–q13 deletion and boys with Fragile X
657 syndrome. *Cognitive Neuropsychology*, 26(2), 172-194. doi:10.1080/02643290802685921
- 658 Woodcock, K. A., Oliver, C., & Humphreys, G. W. (2011). The relationship between specific
659 cognitive impairment and behaviour in Prader-Willi syndrome. *Journal of Intellectual*
660 *Disability Research*, 55, 152-171. doi:10.1111/j.1365-2788.2010.01368.x
- 661 Young, S. E., Friedman, N. P., Miyake, A., Willcutt, E. G., Corley, R. P., Haberstick, B. C.,
662 & Hewitt, J. K. (2009). Behavioral disinhibition: Liability for externalizing spectrum
663 disorders and its genetic and environmental relation to response inhibition across
664 adolescence. *Journal of abnormal psychology*, 118(1), 117-130. doi:10.1037/a0014657
665

666 ***Acknowledgements***

667 The Prader-Willi Associations UK and Ireland supported participant recruitment. Several
668 members of the research team led by the final author assisted with research activities, with
669 particular thanks to Morgan McKenna, Jennifer Norling, and Niamh Rainey. Alex Zylberberg
670 made a major contribution to the design of the neurocognitive tests of switching and
671 associated outcomes. The most important thanks go to the children with PWS and their
672 family members who participated in the research.
673

674 ***Funding statements***

675 The work was supported by a grant awarded to second and corresponding authors by the
676 Foundation for Prader-Willi Research UK and the Foundation for Prader-Willi Research US.

677 **Tables**

678 Table 1. Switching test composite outcomes at each assessment session in the evaluation.
 679 Green shading indicates improvement of at least 5%, orange shading indicates improvement
 680 of 5% or less, red shading indicates worsening of scores of at least 5%.

681

| Participant pseudonym | Mary | Jess | Ellie | May | Sarah | |
|--|--------------------|-------------|--------------|------------|--------------|----------|
| First play phase | Placebo | Active | Placebo | Placebo | Placebo | |
| Age (years) | 13 | 15 | 9 | 10 | 17 | |
| # tests completed at all stages | 3 | 4 | 4 | 3 | 2 | |
| Placebo training (minutes) | 177.5 | 85.83 | 68.61 | 118.01 | 137.31 | |
| Active training (minutes) | 0 | 164.91 | 0 | 0 | 160.3 | |
| Baseline | switch error | 0.28 | 0.19 | 0.41 | 0.39 | 0.29 |
| | switch time (ms) | 1219 | 1263 | 1611 | 928 | 1174 |
| Practise | time from baseline | 24 hours | 22 hours | 14 hours | 17 hours | 24 hours |
| | switch error | 0.19 | 0.14 | 0.24 | 0.38 | 0.29 |
| | switch time (ms) | 1010 | 1276 | 1163 | 680 | 916 |
| | time from baseline | 22 days | 34 days | 35 days | 37 days | 102 days |
| Training phase 1 | switch error | 0.28 | 0.05 | 0.34 | 0.46 | 0.31 |
| | switch time (ms) | 1097 | 1038 | 1434 | 635 | 1170 |
| Training phase 2 | time from baseline | | 198 days | | | 203 days |
| | switch error | | 0.10 | | | 0.31 |
| | switch time (ms) | | 1169 | | | 805 |
| | BRIEF shift | | 12.5% | | | 37.5% |
| Practice effect | error % improved | 30.0 | 27.8 | 41.0 | 3.5 | 0 |
| | time % improved | 17.1 | -1.0 | 27.8 | 26.7 | 22.0 |
| Placebo effect | error % improved | -30.0 | -23.2 | -25.6 | -21.4 | -7.1 |
| | time % improved | -7.1 | -10.4 | -16.9 | 4.9 | -21.7 |
| Active training effect | error % improved | | 44.44 | | | 0 |
| | time % improved | | 18.8 | | | 31.1 |

682

683 **Figure captions**

684 Figure 1. An example of images used in a questionnaire item for children. In this case, the
685 question asked children how much they understood how to play a game (a picture of each
686 game was also provided).

687 Figure 2. An example item used in a questionnaire for children regarding their preferences
688 about existing games (here, regarding their preferences for control systems).

689 Figure 3. Example images used in a three-point Likert-style scale asking children how much
690 they enjoyed playing the first prototype version of the game.

691 Figure 4. Example quiz-style item designed to determine how well children understood what
692 to do in the first prototype version of the game.

693 Figure 5. Gameplay preferences of participating children (pp) with Prader-Willi syndrome.

694 Figure 6. Preferred game control systems of participating children (pp) with Prader-Willi
695 syndrome.

696 Figure 7. An example screen from the prototype game, illustrating the core task switching
697 demand. The player controls the Collector (the character in the center of the screen). Players
698 are required to collect the Creatures of the type indicated in the Target Panel (top left), while
699 avoiding all other Creatures. The core switching demand is provided by changing how the
700 collectible Creatures are identified in the Target Panel (i.e., by their shape or by their colour).
701 In this example, the player must collect cuboidal Creatures and avoid pyramidal Creatures.
702 This image also shows a Power Up (top right), as explained in Supplemental Material. The
703 exclamation point indicates that a Hazard, which the player must avoid, is about to appear at
704 that location (see Supplemental Material).


705 Figure 8. Procedure for pilot evaluation.

706



707
708 *Figure 1*
709
710
711

b. Some of the games were controlled by **TOUCHING** the screen. Some of the games were controlled by **MOVING** the tablet. Which of these did you find easiest to use?*



712
713 *Figure 2*
714
715



716

717


718 *Figure 3*

719

720

721


8 → Look at the picture below. Should they collect the creature?*



722

723

724 *Figure 4*



725


726

727

728 *Figure 5*

729

730



731


732

733 *Figure 6*

734


735

736



737
738
739
740
741

Figure 7



742 T1 assessment → T2 assessment → Training 1 → T3 assessment → Training 2 → T4 assessment

743
744
745
746

Figure 8

Supplemental Material

List of games used in the consultation with children prior to development reported in Study 1.

Websites given are intended to provide the reader with the best possible information about each game. Where possible, an official game website is provided, or a website where a version of the game can be played (all websites last accessed January 31 2019). In the case of Mole Kart, Cordy 2, and Gravity Duck, no official websites or playable versions are available at the time of writing. In these cases, we have provided links to websites which provide the best available information (Wikipedia or YouTube video demonstrating the gameplay).

Tealy and Orangey (<http://www.addictinggames.com/action-games/tealy-and-orangey-game.jsp>)

Multitask (<https://www.kongregate.com/games/icylime/multitask-2>)

Lux Ahoy (<https://luxahoy.com/>)

UFO Run (<http://www.crazygames.com/game/ufo-run>)

Fit it Quick (<https://www.coolmathgames.com/0-fit-it-quick>)

Monument Valley (<https://www.monumentvalleygame.com>)

99 Bricks Wizard Academy (https://en.wikipedia.org/wiki/99_Bricks_Wizard_Academy)

Mole Kart (https://en.wikipedia.org/wiki/Mole_Kart)

Cordy 2 ([https://en.wikipedia.org/wiki/Cordy_\(video_game\)](https://en.wikipedia.org/wiki/Cordy_(video_game)))

Shu's Garden (<http://shusgarden.ca/>)

LEGO Junior's Quest (<https://www.lego.com/en-us/family/apps/quest>)

Dr. Panda Handyman (<https://drpanda.com/games/dr-panda-handyman>)

Amazing Alex (<http://teaser.amazingalex.com/>)

Toca Builders (<https://tocaboca.com/app/toca-builders/>)

Gravity Duck (<https://www.youtube.com/watch?v=O0U-5moIvUk>)

Table S1 Features included in the game

| Feature of the game | Description | Comments |
|-------------------------------|---|--|
| Collector | The character controlled by the player. See Figure 7. | |
| Creatures | The characters which the Collector must collect or avoid. Each creature has a shape and a colour. In all versions of the game discussed here, creatures are either red or blue in colour, and cuboid or pyramidal in shape. Creatures move around the game area, changing direction at random intervals, or when they make contact with the edge of the game area or another creature. See Figures 7 and S1. | |
| Dynamic Difficulty Adjustment | Difficulty is adjusted in three ways: (1) Between games, the difficulty is adjusted by increasing how frequently the target is switched from a colour to a shape or vice versa. At the easiest setting, the target switches after every 6 waves (i.e., only once per game). At the hardest setting, the target switches after every wave. (2) Within games, the difficulty of each wave (see Figure S2 for an explanation of waves) is adjusted by adding or removing creatures (more creatures makes the game harder). (3) Within waves, difficulty is adjusted by adding power-ups (which make the game easier) or hazards (which make the game harder). See Figure S2. | In the placebo version of the game, only difficulty adjustment between waves (i.e., (2)) was used. The active version of the game included all three methods of difficulty adjustment. |
| Ghost Mode | A state of the Collector. When in Ghost Mode, the Collector will pass through non-target creatures and rocks without making contact. The Collector's appearance flickers. Ghost Mode is activated for 3 seconds after the Collector makes contact with a non-collectible Creature, or a Hazard. | Makes the game temporarily easier after the player has made a mistake |

| Feature of the game | Description | Comments |
|-------------------------|--|---|
| Hazard | A rock which appears in a random location (although always a minimum distance from the Collector). The appearance of Hazards is accompanied by an explosion sound effect and a Screen Shake. Hazards pursue the Collector until they either make contact with the Collector, or 3 seconds has elapsed. When either of these conditions is fulfilled, the Hazard disappears. See Figure S1. | This introduces an additional switching demand, as the player must change their goal from attempting to collect Creatures, to avoiding the Hazard. Hazards are introduced when the player has made 2 consecutive successful collections, in order to make the game temporarily more difficult. Hazards also introduce additional required tasks into gameplay, and high levels of concurrent tasks are features of entertainment video games purported to facilitate learning and its transfer. |
| Music and Sound Effects | Music is optional, and can be turned on or off in the game settings. The game also incorporates multiple sound effects. | Some participants' caregivers reported that participants found the music unpleasant; whilst others enjoyed the music. |
| Power Up (fast mode) | A lightning bolt appears in a random location. If the Collector collects the lightning bolt, their velocity is increased for 5 seconds. See Figure 7. | This makes the game temporarily easier, as the Creatures are easier to catch. Power Ups are introduced after the player has made 2 consecutive unsuccessful collection attempts (i.e., they have made contact with creatures that are not currently collectible). |
| Power Up (slow mode) | A clock appears in a random location. If the Collector collects the clock, the velocity of the Creatures is reduced for 5 seconds. See Figure S1. | Provides scaffolding to successful performance. Power Ups also introduce additional required tasks, increasing concurrent task load. |
| Psuedo-3D Graphics | Use of graphical projection to simulate 3 dimensions in 2-dimensional images; also known as 2.5D graphics. See Figure S3. | This entails that the Creatures (which are geometric shapes), appear differently depending on their direction of travel. This introduces an additional switching demand. |
| Scoring | The player receives 1 point for collecting a Creature. If the Collector makes contact with a Creature that is not currently collectible, the player loses 1 point (unless their score is already 0). The current score is displayed in a panel in the bottom left of the screen. See Figure 1. | |
| Screen Shake | The entire contents of the screen move very rapidly in random directions for a moment, as if an earthquake has occurred in the game world. | |

| Feature of the game | Description | Comments |
|---------------------|--|---|
| Target Cue | A representation of the current target (i.e., which colour or shape of creature should be collected) is temporarily displayed in the centre of the screen. | |
| Target Indicator | A white circle which appears around a creature that is currently collectible for 2 seconds. Target Indicators are displayed when the Collector makes contact with a non-collectible Creature and at the beginning of each new wave (see Figure S2 for an explanation of waves). See Figure S1. | Provides scaffolding to the player when the task switches. |
| Target Panel | An image representing the current target (i.e. which colour or shape of creature should be collected), displayed on a black background. The Target Panel can be hidden, in which case only a tab is shown (i.e., the information is not visible; as shown in Figure 2); the player must tap on the tab to reveal the Target Panel, as shown in Figure 1. | The fact that the panel retracts (i.e., the player must tap on it to reveal it) introduces an additional switching demand into the game, in that the player must switch from the current goal (e.g., collection a Creature) to operating the panel. |




Figure S1. A screenshot from the game. The Collector is shown in the center of the screen. Also shown is a Hazard (rock) and a Power Up (Slow Mode; the clock at top right). The player's current score is shown in the score panel (bottom left). In this image, the Target Panel (top left) is shown retracted (cf. Figure 7). Also shown are Target Indicators (i.e., the white circles around some of the creatures). In this example, the player must collect cuboidal Creatures while avoiding pyramidal Creatures.




Figure S2. The concept of waves in the game.




Figure S3. Pseudo-3D graphics used in the game. Each shape can be represented in multiple ways (depending on the direction the creature is moving), simulating a 3D view in 2D graphics.

Table S2 Differences between the placebo and active versions of the game.

| Feature of the game | Active game | Placebo game |
|--|---|---|
| Dynamic Difficulty Adjustment (DDA) | Difficulty adjusted between games, between waves, and within waves. See Table S1 and Figure S2. | Difficulty adjusted between waves only. See Table S1 and Figure S2. |
| Hazards; Power Ups | Included. See Figure S1 | Not included |
| Target Panel | When tapped, displays for 1 second before retracting. See Figures 1 and S1. | Displayed permanently. See Figure 1. |

Practice procedure for switching tests

The four switching tests were each administered following a bespoke practice procedure: Introduction trials introduced the child to the task and checked responses could be linked to target stimuli. Tutorial trials introduced the child to the *cue to task* and checked that the cue could be linked to the correct task. Preparation trials provided the child with an opportunity to prepare for the *measure trials* (would be used to evaluate performance), in which all features were identical to those that would be used to evaluate performance, except that children were provided with feedback contingent on an incorrect response or no response having been provided within the allotted time limit. Thresholds for failure were imposed on each trial type comprised in the practice procedure. Thresholds were selected to strike a balance between giving children the opportunity to demonstrate competence, and maintaining total maximum testing duration acceptably low (see Table S3 for more details).

Switching test characteristics

The four switching tests were designed to each draw to different degrees on the cognitive skills required for appropriate performance, which do not involve task switching. Thus, categorisation decisions ranged from low level perceptual to high level conceptual categories; stimuli were presented to visual and auditory modalities in different tests; *cues to task* were presented to visual and auditory modalities in different tests and additionally indicated the task to different degrees of transparency; and *cues to task* were presented at different durations preceding target stimulus presentation, providing children with different lengths of time for task preparation (see Table S4).

Switching test trial structure

The trial structure differed slightly across introduction, tutorial, preparation and measure trials in order to create the graded practice procedure. However, trial structure was equivalent across all four tests (see Figure S5-S7).

Switching test testing procedure

A storyline about an alien visiting Earth was used to motivate children during engagement with the tests, which involved audio phrases generated by a computer, and images including the well-known alien character from the film *E.T the Extra-Terrestrial* accompanying test explanations and feedback. To allow the tests to be completed flexibly across a range of possible computers at participants' homes, the size of stimuli adapted based on the resolution of the screen being used (which caregivers were instructed to indicate by measuring a line that appeared on the screen following log in). Screen resolutions used varied between 3.20 and 4.60 pixels per millimetre.

Trials administered during the practice procedure were selected so as to best explain what was required to children and ensure the relevant cognitive skills had been tested at each stage. Following practice, the 49 *measure trials* followed the same pre-determined sequence for all tests, with task switches every second trial. Four different *target stimuli* were available for each test, which could either be congruent – when the same response was afforded by both tasks – or non-congruent – when different responses were afforded by each task. Task switches were presented on the third trial and then every second trial. Thus, the first trial was not classified as a switch or a repeat trial. And, from the second trial onwards repeat and switch trials alternated. In this way, trials were balanced for congruency, switching, stimulus and task, with three trials of each combination of these features.

Table S3: Description of practice procedure for switching tests

| Trial type & # available | Trial function: to assess... | Repeat procedure | Threshold for failure | Scaffolding | Procedure (also see figures S5-9) |
|-------------------------------------|--|--|--|--|---|
| Introduction (IT) 4 | understanding of the two tasks | Correct response followed by next trial in sequence; | Incorrect response or time out for any single trial 5 times | Verbal explanation of task and response mapping; verbal and visual feedback on success (correct/ incorrect/ time out) | 1. <i>Target stimulus</i> at top or bottom centre + <i>cue to response</i> presented until response or 5s (time out); 2. Trial feedback presented for duration of verbal feedback sound |
| Tutorial (TT) 4 | understanding of the task cues; and that task switches can occur | Incorrect response or no response before time out, followed by a repeat of the same trial | Incorrect response or time out for any single trial 3 times | Verbal explanation of task cue; verbal and visual feedback on success | 1. <i>Cue to task</i> + verbal description of cue, presented for duration of verbal description (longer for trials 1 & 3, see script); 2. Addition of <i>Target stimulus</i> + <i>cue to response</i> (as ITs) ; 3. Trial feedback (as ITs) |
| Preparation (PT) 4 | ability to task switch in the context of the test | Any response is followed by next trial in sequence; If > 1 incorrect response or no response before time out, following 4 th trial all 4 trials are repeated in sequence | At least 2 incorrect or time out responses in the 3 rd repetition of all 4 trials | Reminder of task cue on trials 1 and 3 only; verbal and visual feedback if response incorrect or too slow; encouragement feedback following trial 4 if trials to be repeated | As TTs except: a. time out is 3s; b. reduced verbal description of cue (see audio script) and only on trials 1 & 3; c. no trial feedback presented following correct responses |

Table S4: Description of tasks, required responses and stimuli for switching tests.

| Test | Type of discrimination | Tasks 1 | Cue to task | Task preparation | Response mapping | Cue to response | Target stimuli |
|---------------|---|---|--|--|--|--|---|
| Location size | Conceptual category (relatively uncommon) | 1. Is stimulus usually inside or outside the house? 2. Will stimulus fit inside a rucksack or not? | Pictorial (transparent) A: house cartoon B: rucksack cartoon | Relatively long (cue presented 1000ms before target) | Left: in Right: out | Semi-transparent; High response conflict; L: bottom L side in symbol R: bottom R side out symbol | Verbal 1. “Toaster”; 2. “Donkey”; 3. “Bookshelf”; 4. “Football” |
| Age gender | Conceptual category (relatively common) | 1. Is stimulus young or old? 2. Is stimulus male or female? | Locational; High response conflict A: top L or R B: bottom L or R | Shortest (cue presented concurrently with target) | L: young/female R: old/male | Semi-transparent; For response conflict, see <i>Cue to task</i> L: top children’s sign & bottom men’s sign to slight L of centre R: top old person’s sign & bottom women’s sign to slight R of centre | Pictorial 1. Boy 2. Old woman 3. Girl 4. Old man |
| Shape colour | Perceptual category (low level) | 1. Is stimulus a square or a circle? 2. Is stimulus red or blue? | Verbal A: “shape” B: “colour” | Longest (cue presented 2000ms before target) | Left: square/red Right: circle/blue | Transparent; High response conflict L: bottom left side red square R: bottom right side blue circle | Pictorial 1. Red square 2. Blue circle 3. Blue square 4. Red circle |

| | | | | | | | |
|--------------|------------------------------------|---|-----------------------------|--|---------------------------------|--|---|
| Global local | Perceptual category (higher level) | 1. Is global shape a square or a triangle? 2. Is local shape a square or a triangle? | Pictorial (non-transparent) | Relatively short (cue presented 500ms before target) | Left: square Right: triangle | Transparent; For response conflict, see <i>Stimuli</i> L: bottom left side black outline of square R: bottom right side black outline of triangle | Pictorial (Navon) High response conflict 1. Square of squares 2. Triangle of triangles 3. Triangle of squares 4. Square of triangles |
|--------------|------------------------------------|---|-----------------------------|--|---------------------------------|--|---|




Figure S5: Summary of switching test structure

| | | | | |
|---------------------------|---|--|---|--|
| [moon image] | [house image] | “donkey” | A picture of ET in front of an untidy room | A picture of a satchel with a picture of ET on the front |
| Fixation (PTs & MTs only) | Cue to task (not in ITs) | Target stimulus + cue to response | Trial feedback (not following correct PTs or any MTs) | Encouragement feedback (only following every 4 PTs; and 7 MTs) |
| PTs: 500ms; MTs: 300ms | TTs/PTs: sound duration or 2000ms if no sound; MTs: test specific | Until response or timeout IT/TT timeout= 5s PT/MT timeout=3s | Sound duration | Sound duration |

Figure S6: Summary of trial structure for switching tests: The fixation image was the same for all tests. Stimuli from the Location Size test; on a location task trial; and assuming an incorrect or time out response; are used to illustrate other trial components. Dotted lines indicate trial components that are not comprised in all trial types.

Switching test removal of outliers

For each switching test, switch time and switch error outcome variables were calculated based on the 24 *switching* trials that followed a task switch. Prior to calculation of switch time outcomes, trials were examined for those with reaction times lying outside three standard deviations unit from the participant's mean reaction time for switching trials in the relevant task. However, no such outlying trials were identified.

Supplementary evaluation results

Switch time and switch error practice effects were calculated based on the percentage improvement in scores between T1 and T2 assessments. Improvement in these scores linked to training phase 1 was calculated based on the percentage improvement between T1 and T3 assessments, with practice effects subtracted from this value. In the corresponding manner, improvement linked to training phase 2 was calculated based on the percentage improvement between T1 and T4 assessments, with practice effects subtracted from this value.